

Victor B. Rybakov,\* Eugene V.  
Babaev and Ylya V. DlinnykhChemistry Department, Moscow State Univer-  
sity, 119899 Moscow, RussiaCorrespondence e-mail:  
rybakov@biocryst.phys.msu.su

## Key indicators


Single-crystal X-ray study  
 $T = 293$  K  
Mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å  
 $R$  factor = 0.047  
 $wR$  factor = 0.117  
Data-to-parameter ratio = 9.7For details of how these key indicators were  
automatically derived from the article, see  
<http://journals.iucr.org/e>. $N^S, N^O$ -Diphenyl(thiooxamide)

The molecule of the title compound,  $\text{PhNHC}(=\text{S})-\text{C}(=\text{O})\text{NHPh}$  or  $\text{C}_{14}\text{H}_{12}\text{N}_2\text{OS}$ , consists of two planar fragments, one of which includes the central  $\text{NHC}(=\text{S})-\text{C}(=\text{O})\text{NH}$  chain together with the Ph substituent on the  $\text{C}=\text{O}$  side; the second Ph ring, the one on the  $\text{C}=\text{S}$  side, all by itself, makes up the second planar fragment. Its plane is twisted about the  $\text{N}-\text{C}$  bond by  $52.87$  ( $9$ ) $^\circ$  with respect to the plane of the first fragment. The  $\text{C}=\text{S}$  and  $\text{C}=\text{O}$  double bonds adopt the *transoid* conformation with the torsion angle  $\text{S}=\text{C}-\text{C}=\text{O}$  equal to  $-179.6$  ( $2$ ) $^\circ$ . The molecules in the crystal are linked into centrosymmetric dimers due to the  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bond involving the thioamide NH group.

Received 8 May 2001  
Accepted 27 July 2001  
Online 31 July 2001

## Comment

The molecular structure of the title compound is shown in Fig. 1. The molecule consists of two planar fragments: the  $\text{C}_9-\text{C}_{14}$  phenyl ring makes up one of them and the  $\text{C}_3-\text{C}_8$  phenyl ring together with the  $\text{N}_1-\text{C}_1(\text{O}_1)-\text{C}_2(\text{S}_1)-\text{N}_2$  chain atoms attached to  $\text{C}_3$  forms the other one. The intermolecular hydrogen bond [ $\text{N}_2\cdots\text{O}_1^i$  2.42 (3) Å,  $\text{N}_2\cdots\text{O}_1^i$  3.136 (4) Å and  $\text{N}_2-\text{H}_2\cdots\text{O}_1^i$  148 (3) $^\circ$ ; symmetry code: (i)  $1-x, 1-y, -z$ ] links the molecules in the crystal into centrosymmetric dimers. In the thiooxamide part of the molecule, the S and O atoms are *trans* with respect to each other; the



torsion angle  $\text{S}_1-\text{C}_2-\text{C}_1-\text{O}_1$  is  $-179.6$  ( $2$ ) $^\circ$ . The conjugation between the thioamide and amide moieties of the molecule is weak, as the  $\text{C}_1-\text{C}_2$  bond [1.532 (4) Å] is significantly longer than the standard  $\text{Csp}^2-\text{Csp}^2$  bond length in conjugated systems (1.46–1.48 Å; Allen *et al.*, 1987). The bond distances  $\text{C}_1=\text{O}_1$  [1.231 (3) Å] and  $\text{N}_1-\text{C}_3$  [1.413 (4) Å] are longer than the standard bond lengths for  $\text{C}=\text{O}$  (1.22 Å) and  $\text{C}_{\text{Ar}}-\text{Nsp}^2$  (1.36 Å) (Allen *et al.*, 1987). This effect may be explained by a weak delocalization of electron density in the amide group. The bond angle  $\text{C}_1-\text{N}_1-\text{C}_3$  has an abnormal value of  $131.5$  ( $3$ ) $^\circ$ . This large bond angle may be explained by steric strain in the planar fragment  $\text{O}_1-\text{C}_1-\text{N}_1-\text{C}_3-\text{C}_4-\text{H}_4$ . The second phenyl ring ( $\text{C}_9-\text{C}_{14}$ ) is twisted about the  $\text{N}_2-\text{C}_9$  bond with respect to the planar moiety  $\text{N}_2-\text{C}_2(\text{S}_1)-\text{C}_1(\text{O}_1)-\text{N}_1-(\text{C}_3-\text{C}_8)$  by  $52.87$  ( $9$ ) $^\circ$ . Due to this twisting, the  $\text{C}_2-\text{N}_2-\text{C}_9$  bond angle is not distorted and has a generally accepted value of  $126.1$  ( $3$ ) $^\circ$ . Only one related structure

(Krayushkin *et al.*, 1996) (with *n*-butyl instead of phenyl as in our case) was found in the Cambridge Structural Database (Allen & Kennard, 1993). The main structural features of this molecule are essentially identical with those of the title compound.

## Experimental

A mixture of 0.5 g (2 mmol) 3-benzoxazolo[3,2-*a*]pyridinium-2-olate and 10 ml thionyl chloride (SOCl<sub>2</sub>) was kept under reflux for 1 h. Thionyl chloride was evaporated and the precipitate obtained was washed with benzene (3 × 10 ml) and dissolved in dichloromethane CH<sub>2</sub>Cl<sub>2</sub> (20 ml). 0.4 g (4.3 mmol) aniline (C<sub>6</sub>H<sub>7</sub>N) was added to this solution. After heating at 313 K for 1 h and leaving to stand overnight at room temperature, the dichloromethane was evaporated, and the residue washed with water (3 × 20 ml) and recrystallized from a mixture of chloroform and diethyl ether (4:1). The yield was 0.2 g (37%).

### Crystal data

C <sub>14</sub> H <sub>12</sub> N <sub>2</sub> OS	$D_x = 1.390 \text{ Mg m}^{-3}$
$M_r = 256.32$	Mo $K\alpha$ radiation
Monoclinic, $P2_1/c$	Cell parameters from 25 reflections
$a = 4.026 (5) \text{ \AA}$	$\theta = 13.0\text{--}15.0^\circ$
$b = 14.682 (9) \text{ \AA}$	$\mu = 0.25 \text{ mm}^{-1}$
$c = 20.728 (12) \text{ \AA}$	$T = 293 (2) \text{ K}$
$\beta = 90.27 (2)^\circ$	Prism, yellow
$V = 1225.1 (17) \text{ \AA}^3$	$0.38 \times 0.12 \times 0.06 \text{ mm}$
$Z = 4$	

### Data collection


Enraf–Nonius CAD-4 diffractometer	$h = -4 \rightarrow 4$
$\omega$ scans	$k = 0 \rightarrow 17$
2089 measured reflections	$l = 0 \rightarrow 24$
2051 independent reflections	2 standard reflections every 200 reflections
1287 reflections with $I > 2\sigma(I)$	frequency: 60 min
$R_{\text{int}} = 0.070$	intensity decay: none
$\theta_{\text{max}} = 25.0^\circ$	

### Refinement

Refinement on $F^2$	All H-atom parameters refined
$R[F^2 > 2\sigma(F^2)] = 0.047$	$w = 1/[\sigma^2(F_o^2) + (0.0597P)^2]$
$wR(F^2) = 0.117$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.95$	$(\Delta/\sigma)_{\text{max}} = 0.048$
2051 reflections	$\Delta\rho_{\text{max}} = 0.20 \text{ e \AA}^{-3}$
212 parameters	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$

**Table 1**  
Selected geometric parameters ( $\text{\AA}$ ,  $^\circ$ ).

S1–C2	1.660 (3)	N1–H1	0.85 (3)
O1–C1	1.231 (3)	C2–N2	1.310 (4)
C1–N1	1.316 (4)	N2–C9	1.428 (4)
C1–C2	1.532 (4)	N2–H2	0.81 (3)
N1–C3	1.413 (4)		
O1–C1–N1	126.9 (3)	C1–C2–S1	121.6 (2)
O1–C1–C2	119.3 (3)	C2–N2–C9	126.1 (3)
N1–C1–C2	113.7 (2)	C4–C3–N1	123.0 (3)
C1–N1–C3	131.5 (3)	C8–C3–N1	117.0 (3)
N2–C2–C1	112.1 (2)	C14–C9–N2	120.2 (3)
N2–C2–S1	126.3 (2)	C10–C9–N2	119.2 (3)



**Figure 1**  
ORTEP-3 (Farrugia, 1998) view of the molecule of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.

**Table 2**  
Hydrogen-bonding geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D\text{--}H\cdots A$	$D\text{--}H$	$H\cdots A$	$D\cdots A$	$D\text{--}H\cdots A$
N1–H1 $\cdots$ S1	0.85 (3)	2.38 (3)	2.938 (3)	124 (3)
N2–H2 $\cdots$ O1	0.81 (3)	2.17 (4)	2.632 (4)	116 (3)
C4–H4 $\cdots$ O1	0.88 (3)	2.45 (3)	2.977 (4)	119 (3)
N2–H2 $\cdots$ O1 <sup>i</sup>	0.81 (3)	2.42 (3)	3.136 (4)	148 (3)
C14–H14 $\cdots$ S1	0.93 (3)	2.97 (3)	3.250 (4)	99 (2)

Symmetry code: (i)  $1 - x, 1 - y, -z$ .

All H atoms were refined isotropically; the C–H bonds are in the range 0.82–1.00  $\text{\AA}$ .

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *WinGX98* (Farrugia, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3* (Farrugia, 1998).

This work has been supported by the Russian Foundation for Basic Research (project No. 99–03–33076). We also acknowledge the support of this Foundation in payment of the licence for using the Cambridge Structural Database (project No. 99–07–90133).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Allen, F. H. & Kennard, O. (1993). *Chem. Des. Autom. News*, **8**, 1, 31–37.
- Enraf–Nonius (1989). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1998). *ORTEP-3 for Windows and WinGX98*. University of Glasgow, Scotland.
- Krayushkin, M. M., Vorontsova, L. G., Kurella, M. G., Zavarzin, I. V. & Yarovenko, V. N. (1996). *Russ. Chem. Bull.* **2**, 485–487.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.