

## Efficient synthesis of 5-substituted 2-aryl-6-cyanoindolizines via nucleophilic substitution reactions

Eugene V Babaev\*, Natalya I Vasilevich and Anna S Ivushkina

Address: Department of Chemistry, Moscow State University, 119992, Moscow, Russia

Email: Eugene V Babaev\* - babaev@org.chem.msu.su; Natalya I Vasilevich - nikto25@hotmail.com; Anna S Ivushkina - anna@hotmail.ru

\* Corresponding author

Published: 07 October 2005

Received: 11 June 2005

Beilstein Journal of Organic Chemistry 2005, 1:9 doi:10.1186/1860-5397-1-9

Accepted: 07 October 2005

This article is available from: <http://bjoc.beilstein-journals.org/content/1/1/9>

© 2005 Babaev et al; licensee Beilstein-Institut.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract


2-Aryl-6-cyano-7-methyl-5-indolizines were successfully converted into 2-aryl-5-chloro-6-cyano-7-methylindolizines. The obtained 5-chloroindolizines readily underwent nucleophilic substitution at position 5 leading in high yields to novel 5-functionalised indolizines.

Indolizines are an important class of heterocyclic compounds since many natural alkaloids contain in their structure a saturated (swainsonine) or aromatic (campothecin) indolizine moiety. While the chemistry of indolizines has been widely investigated[1] the chemistry of 5-substituted indolizines remains very poor because there are only a few reliable ways for their synthesis.

8-Nitroindolizines may undergo amination at position 5 ( $S_NH$  substitution) under the action of secondary amines.[2] 2-Phenylindolizine can be lithiated at position 5, and the resulting indolizyl lithium can react with some electrophiles ( $CO_2$ , PhCHO, PhCN,  $Me_3SiCl$ , MeI) leading to variety of new products in good yields.[3] An interesting method for preparing 5-substituted indolizines by recyclization of oxazolo[3,2-a]pyridinium salts was developed in our laboratory.[4,5] Using this strategy a series of 5-substituted indolizines have been prepared in good yields, but (although the method seems to be quite reliable) it is currently restricted only by secondary amines.

In seeking for a synthetic approach to 5-substituted indolizines we have assumed that indolizines bearing an appropriate leaving group (*e.g.* halogen) at position 5 may undergo nucleophilic substitution. Herein we discuss the synthesis of previously unknown 5-chloroindolizines and their use as precursors to novel 5-substituted indolizines via nucleophilic displacement reactions.

The synthesis of 2-aryl-5-chloro-6-cyano-7-methylindolizines **2** is shown in Scheme 1. 2-Aryl-6-cyano-7-methyl-5-indolizines **1 a – d** were prepared according to protocol of Gevald.[6] Our modification of the original method included separation of N- and O-isomers of phenacyl pyridines before cyclization (using the difference in their solubility in chloroform). Although  $^1H$ -NMR (see Additional file 2) and Nuclear Overhauser Effect confirmed the structure **A** for indolizines **1**, we assumed the existence of tautomerism between forms **A** and **B** involving hydrogen interchange between oxygen and C-3 carbon (Scheme 1). Although the amount of tautomer **B** is negligibly small, one would expect that treatment of



**Scheme 1:** Synthesis of 2-aryl-5-chloro-6-cyano-7-methylindolizines **2**. Possible tautomeric structures **A** and **B** for 2-aryl-6-cyano-7-methyl-5-indolizinones **1**.

**1 a - d** with phosphorous oxychloride may lead to substitution of oxo/oxy-group to chlorine giving the products **2 a - d**. (It is well known that analogous 2-hydroxypyridines which exist in the pyridone tautomeric form can be easily converted to 2-chloroderivatives by reaction with POCl<sub>3</sub>). [7] Indeed, heating of indolizinones **1 a - d** in POCl<sub>3</sub> at 80–100°C during 10 hours without any solvent followed by pouring into a mixture of ice/sodium acetate and filtration of the green precipitate afforded crude 5-chloroindolizines. After column chromatography (eluent – carbon tetrachloride) yellow solids were obtained. Performing this reaction in the presence of two-fold molar excess of trimethylbenzylammonium chloride or TEBAC increased the yields of **2 a - d** up to 30–75%. In the <sup>1</sup>H NMR spectra of these products<sup>†</sup> the initial signal of 3-CH<sub>2</sub> group at ~5 ppm (intensity 2H) disappeared, and a new aromatic signal 3-CH (with intensity 1H) appeared at 7.99 – 8.11 ppm.

The halogen atom in 5-chloro-6-cyanoindolizines **2** should be activated to nucleophilic substitution reactions by the suitable *ortho*-arrangement of the nitrogen atom of the pyridine ring and electron-withdrawing cyano-group. The pattern strongly resembled 2-chloro-3-cyanopyridine,

that is why we anticipated successful substitution in reactions of **2** with oxygen, nitrogen, and sulfur nucleophiles. Indeed, 5-chloroindolizines readily underwent nucleophilic substitution to produce previously unknown compounds **3 - 6** in good to excellent yields (Scheme 2). These products are detailed in Table 2. Thus, 5-methoxyindolizines **3 a - c** were formed after refluxing **2 b - d** in solution of sodium methoxide in methanol overnight in good yields (Scheme 2). Treatment of **2a, d** with excess of amines without any solvent gave 5-amino derivatives **4 a - h**. In the case of secondary amines (**4 a - c, e - g**) the reaction proceeded at room temperature, but reaction with less nucleophilic benzylamine (**4 d, h**) required heating for 30 min. Nucleophilic substitution also occurred with sulfur nucleophiles. Thus, **2d** reacted with mercaptoethanol under basic conditions leading to **5a**. Conversion **2a** into **5b** was conveniently achieved with ethyl mercaptoacetate in ethanolic sodium hydroxide. Interestingly, **2a** reacted also with thiourea in refluxing butanol giving indolizine-thione **6**. The product **6** seems to be the result of decomposition of unstable isothiuronium salt, and the process resembles the known conversion of 2-chloro-3-cyanopyridines to 3-cyanopyridine-thiones under the same conditions via a similar intermedi-


ate. [8] <sup>1</sup>H NMR spectrum of **6**, which was very similar to the spectra of **1**, indicated disappearance of aromatic proton signal H<sub>3</sub> and appearance of a signal at 5.35 ppm with intensity 2H.†

In conclusion, we are the first to obtain 2-aryl-5-chloro-6-cyano-7-methylindolizines from 2-aryl-6-cyano-7-methyl-5-indolizinones and to prove the possibility to employ them in nucleophilic substitution reactions. Moreover, these reactions are the first examples of preparative nucleophilic substitution in indolizines, and our findings open a new way to functionalize the C-5 position (in most cases considered as inactive). The studies of further cyclizations of 5-substituted indolizines involving neighbouring cyano-group and ring position C<sub>3</sub> is underway.

### Note

\*Characteristics of parent indolizinones **1 a – d** were identical to those described in literature.[9]

† Supporting information



**Scheme 2: Nucleophilic substitution in 2-aryl-5-chloro-6-cyano-7-methylindolizines.**

**Table 1: Properties of 5-substituted 2-aryl-6-cyano-7-methylindolizines**

No.	5-X*	R in 2-Ar	Yield %	m.p., °C
<b>2 a</b>	Cl	p-F	30	173–175
<b>2 b</b>	Cl	p-Cl	65	198–200
<b>2 c</b>	Cl	p-Br	54	229–230
<b>2 d</b>	Cl	p-Me	74	157–158
<b>3 a</b>	OMe	p-Cl	50	169–172
<b>3 b</b>	OMe	p-Br	71	197–200
<b>3 c</b>	OMe	p-Me	73	170–173
<b>4 a</b>	pyrrolidyl	p-F	99	189–190
<b>4 b</b>	piperidyl	p-F	91	205–209
<b>4 c</b>	hexamethylenimino	p-F	88	186–188
<b>4 d</b>	benzylamino	p-F	58	190–194
<b>4 e</b>	pyrrolidyl	p-Me	74	228–230
<b>4 f</b>	piperidyl	p-Me	71	212–215
<b>4 g</b>	hexamethylenimino	p-Me	85	213–216
<b>4 h</b>	benzylamino	p-Me	83	185–187
<b>5 a</b>	S(CH <sub>2</sub> ) <sub>2</sub> OH	p-Me	71	132–135
<b>5 b</b>	SCH <sub>2</sub> CO <sub>2</sub> Et	p-F	70	142–145

### Additional material

#### Additional File 1

Supporting tables

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1860-5397-1-9-S1.doc>]

#### Additional File 2

Supporting information

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1860-5397-1-9-S2.pdf>]

### Acknowledgements

This work was supported by Russian Foundation of Basic Research (grant 04-03-32823).

### References

- Fitch WV: Pyrroles with fused six-membered heterocyclic rings: a-fused. In *Comprehensive Heterocyclic Chemistry Volume 4*. Edited by: Katritzky AR, Rees CW. London: Pergamon Press; 1984:443-478.
- Kost AN, Sagitullin RS, Gromov SP: *Heterocycles* 1977, **7**:997-1001.
- Renard M, Gubin J: *Tetrahedron Lett* 1992, **33**:4433-4434.
- Babaev EV, Efimov AV, Maiboroda DA, Jug K: *Eur J Org Chem* 1998, **1**:193-196.
- Babaev EV: *J Heterocycl Chem* 2000, **37**:519-526.
- Gevald K, Jansch HJ: *J Prakt Chem* 1976, **318**:313-320.
- Kwok R: *J Heterocycl Chem* 1978, **15**:877-880.
- Kitaygorodova EA, Konyushkin LD, Mikhaychenko SN: *Khim Geterotsikl Soedin* 1999, **3**:337-341. (Rus)
- Lin CF, Lin YF, Lo YC, Chen KT, Su TL: *Heterocycles* 2000, **1**:15-26.